



Eksportpakke – Læggekartofler sæson 2025/2026

Indhold

NYT SIDEN SIDST	2
ALBANIEN	3
ALGERIET	4
ARMENIEN	6
ASERBAJDSJAN	7
BOSNIEN-HERCEGOVINA	8
EGYPTEN	9
FORENEDE ARABISKE EMIRATER	11
IRAK – MODEL A	12
IRAK – MODEL B	14
ISLAND	16
ISRAEL	17
JORDAN – MODEL A	20
JORDAN – MODEL B	22
KANARISKE ØER	24
KOSOVO	25
LA RÉUNION	26
LIBANON	27
LIBYEN	29
MAROKKO	31
MONTENEGRO	32
NORDMAKEDONIEN	33
SAUDI ARABIEN – MODEL A	34
SAUDI ARABIEN – MODEL B	35
SERBIEN	36
SYRIEN	37
TUNESIEN	38
TYRKIET	39
UK – MERISTEM	40
UK – EKSPORT TIL AVLER	41
UKRAINE	42

Senest gennemgået d. 24-09-2025



Nyt siden sidst

Ved eksport til lande der ikke fremgår af eksportpakken, bedes I kontakte Sundhedfroel@lfst.dk for eksportbetingelser. Følgende lande er fjernet fra listen i 2025:

- Kasakhstan
- Nicaragua
- Pakistan
- Qatar
- Senegal
- Sudan



Albanien

Oplysninger af ukendt oprindelse.

Marksyn

Partikontrol

Visuelt syn:

- 0 % *Phoma* spp.
- 0 % Pulverskurv (*Spongospora subterranea*) (knolde med åbne pustler)
- *Meloidogyne* spp.

Frihed for *Leptinotarsa decemlineata* og *Ditylenchus dipsaci*

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være testet for ringbakteriose

Tillægserklæring på plantesundhedscertifikat

The seed potatoes were grown in land in which Potato Wart Disease (*Synchytrium endobioticum*) has not been known to occur at any time nor within 10 kilometers thereof.

The seed potatoes originate from land which was officially sampled in [redacted] år and examined for the presence of commercially harmful nematodes which would be subject to quarantine regulations. Soil samples were found to be free of the potato cyst nematodes *Globodera pallida* and *Globodera rostochiensis*. The official inspection of the crops during the period of growth did not reveal any evidence of the presence of potato cyst nematodes.

Potato tuber moth (*Phthorimaea operculella*), Japanese beetle (*Popillia japonica*), Potato brown rot (*Ralstonia solanacearum*), Potato Stolbur phytoplasma, Potato yellowing vein 'virus', Potato yellow dwarf virus (PYDV) and Potato Spindle Tuber Viroïd (PSTVd) has never been found on potato in Denmark.

Colorado beetles (*Leptinotarsa decemlineata*), Potato rot nematode (*Ditylenchus destructor*), Potato ring rot (*Clavibacter michiganensis* subsp. *sepedonicus*) and Tomato spotted wilt virus (TSWV) are known not to occur in the consignment.

The crops from which the seed potatoes originate were officially inspected whilst growing, by Inspectors of the Ministry and were certified to conform to the standards of freedom of virus diseases according to the Ministry's certification scheme.

The consignment has been inspected and found substantially free from Gangrene (*Phoma exigua* var. *foveata*), Powdery scab (*Spongospora subterranea*), *Meloidogyne chitwoodi* and *Meloidogyne fallax*.

Øvrigt

Quality certificate skal vedlægges plantesundhedscertifikatet



Algeriet

I henhold til: Autorisation technique d'importation du matériel vegetal,
N124/DPVCT/SDCT/2016

Marksyn

Partikontrol

Der skal foretages et syn af hvert høstnummer til sikring af frihed for visuelle tegn på forekomst af:

- *Phoma exigua* var. *foveata*
- *Phoma indina*
- *Clavibacter michiganensis* subsp. *sepedonicus*
- *Ralstonia solanacearum*
- *Globodera pallida*
- *Globodera rostochiensis*
- *Ditylenchus destructor*
- Potato spindle tuber viroid
- Potato T. trichovirus
- Potato Andean latent virus
- Potato Andean mottle comovirus
- *Synchytrium endobioticum*
- Tomato spotted wilt tospovirus
- *Phytoplasma solani*
- *Boeremia foveata*
- *Ditylenchus dipsaci*
- *Meloidogyne* sp.

Der skal være synet 250 hkg for hvert høstnummer. I de tilfælde, hvor den aflede mængde udgør mindre end 250 hkg, synes den anmeldte mængde.

Hvis der findes tegn på forekomst af kraterråd skal partiet afvises til eksport til Algeriet, medmindre det omsorteres først.

Laboratorieundersøgelser

Sorten skal være testet for ring- og brunbakteriose.

Tillægserklæring plantesundhedscertifikat

The seed potatoes are free from Potato Wart Disease (*Synchytrium endobioticum*).

The seed potatoes originate from land which was officially sampled in [redacted] and examined for the presence of commercially harmful nematodes which would be subject to quarantine regulations. Soil samples were found to be free of the potato cyst nematodes *Globodera pallida* and *Globodera rostochiensis*. The official inspection of the crops during the period of growth did not reveal any evidence of the presence of potato cyst nematodes.

Potato tuber moth (*Phthorimaea operculella*), Japanese beetle (*Popillia japonica*), *Meloidogyne chitwoodi*, *Meloidogyne fallax*, Potato brown rot (*Ralstonia solanacearum*), Potato Stolbur phytoplasma, *Phoma exigua* var. *foveata*, *Phoma andina*, Potato yellowing vein 'virus', Potato yellow dwarf virus (PYDV), Potato Spindle Tuber Viroïd (PSTVd), Potato



T. trichovirus, Potato Andean latentvirus, and Potato Andean mottle comovirus has never been found on potatoes in Denmark.

Colorado beetles (*Leptinotarsa decemlineata*), Potato rot nematode (*Ditylenchus destructor*), Potato ring rot (*Clavibacter michiganensis* subsp. *sepedonicus*) and Tomato spotted wilt virus (TSWV) are known not to occur in the consignment.

The crops from which the seed potatoes originate were officially inspected whilst growing, by Inspectors of the Ministry and were certified to conform to the standards of freedom of virus diseases according to the Ministry's certification scheme.

The consignment has been inspected and found substantially free from *Meloidogyne chitwoodi* and *Meloidogyne fallax*.

The Potato tubers are free from:

- Bacterial wilt (*Clavibacter michiganensis* subsp. *sepedonicus*) and come from crops which have undergone official inspection during the growing season.
- Wart gall (*Synchytrium endobioticum* (schilb. Perc.))
- Phoma (*Phoma exigua* var. *foveata*, *Phoma andina* and *Boeremia foveata*)
- Viable cysts of golden nematodes (*Globodera pallida* and *Globodera rostochiensis*) and come from fields known to be free from these parasites.
- *Ditylenchus dipsaci*
- Potato spindle tuber viroid, Potato T. trichovirus, Potato Andean mottle comovirus and come from countries free from this disease.

Øvrigt

Følgende skal vedlægges plantesundhedscertifikatet:

[Frihedserklæringen for GMO findes på Landbrugsstyrelsens hjemmesiden](#)

Importtilladelse



Armenien

I henhold til: council of Eurasian Economic Commission no 157 og amendment No. 109 af 15. Juli 2022.

Marksyn

Partikontrol

Praktisk taget fri for jord (max 1 %)

Produktionsstedet skal være fri for:

- *Ralstonia solanacearum*
- *Meloidogyne chitwoodi*
- *Meloidogyne fallax*
- *Synchytrium endobioticum*

Frihed for:

- Potato spindle tuber viroid
- *Candidatus liberibacter solanacearum* (zebra chip)

Laboratorieundersøgelser

- *Globodera pallida*
- *Globodera rostochiensis*

Tillægserklæring på plantesundhedscertifikat

Regulated products are grown in zones, places and (or) sites of production free from quarantine pests in accordance with Common Requirements approved by the Decision No. 157 of the Council of the Eurasian Economic Commission of November 30. 2016 and amendment No. 109 of July 15. 2022.

Øvrigt



Aserbajdsjan

I henhold til Phytosanitary norms and rules on regulated articles imported into and transited through the territory of the Republic of Azerbaijan, No. 41, 26 december 2023

Marksyn

Partikontrol

Phoma foveata

Meloidogyne chitwoodi

Meloidogyne fallax

Potato spindle tuber viroid

Candidatus liberibacter solanacearum

Synchytrium endobioticum

Maksimum jord 1 pct. Af den samlede vægt, dog 0,1 pct. hvis *Ralstonia solanacearum* og *Globodera rostochiensis* findes i landet

Laboratorieundersøgelser

Globodera pallida

Globodera rostochiensis

Ralstonia solanacearum

Clavibacter michiganensis subsp. *sepedonicus*

Tillægserklæring på plantesundhedscertifikat

The tubers are free from *Tecia solanivora*, *Epitrix cucumeris*, *Epitrix tuberis*, *Phthorimaea operculella*, Potato Andean latent tymovirus, *Phoma foveata*, *Premnotrypes* spp., *Ditylenchus dipsaci*, *Ditylenchus destructor*, *Globodera pallida*, *Globodera rostochiensis*, *Meloidogyne chitwoodi*, *Meloidogyne fallax*, *Ralstonia solanacearum*, *Clavibacter michiganensis* subsp. *sepedonicus*, *Xiphinema rivesi*, Potato spindle tuber viroid, *Candidatus liberibacter solanacearum*, Tomato spotted wilt virus, *Synchytrium endobioticum*, *Thecaphora solani*.

Øvrigt



Bosnien-Hercegovina

Oplysninger fra: Rulebook on lists of harmful organisms, lists of plants, plant products and regulated objects

Marksyn

Partikontrol

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være testet for ringbakteriose.

Tillægserklæring på plantesundhedscertifikat

Consignment complies with Annex IV.A.I, points 25.1 option (a); 25.2 option (a) and 25.4 option (a), (b), (c) and (dd); 25.4.2 (a) of Book of Rules on lists of harmful organisms (Official Gazette BiH, 48/13).

The seed potatoes were grown in areas where potato wart disease (*Synchytrium endobioticum*) are not known to occur and no symptoms of potato wart disease has been observed either at the place of production or in its immediate vicinity (10 km).

Soil has been tested and found free from the potato cyst nematodes *Globodera pallida* and *Globodera rostochiensis*.

Denmark is known to be free from potato ring rot (*Clavibacter michiganensis* subsp. *sepedonicus*).

Potato brown rot (*Ralstonia solanacearum*, syn. *Pseudomonas solanacearum*) has never been found in Denmark.

Guatemalan potato moth (*Tecia solanivora*, syn. *Scrobipalopsis solanivora*), and Potato Stolbur phytoplasma have never been found in Denmark.

The tubers come from an area free from *Meloidogyne chitwoodi* and *Meloidogyne fallax*.

The crops from which the seed potatoes originate were officially inspected whilst growing, by Inspectors of the Ministry and were certified to conform to the standards of freedom of virus diseases and quarantine diseases according to the Ministry's certification scheme.

The variety purity is 99,9 %. Free from virus diseases in accordance with the official requirements for the class .

Øvrigt

Kartoffelbrok (*Synchytrium endobioticum*) må ikke have været fundet inden for en radius af 10 kilometer af marken kartoflerne stammer fra.



Egypten

Jf. Decree No. 748 of 2019 of the Egyptian Ministry of Agriculture and annex no. 14 of the Ministril Decree no. 562/2019

Marksyn

Godkendt som min. klasse E.

Der må ikke være forekomst af *Dickeya*, *Pectobacterium carotovorum*, *P. atrosepticum* på produktionsarealet i indeværende år og de to foregående to år.

Partikontrol

Der skal foretages et rekvireret syn af hvert høstnummer til sikring af de Egyptiske krav. Der skal være synet minimum 250 hkg pr. høstnummer.

Der synes jf. Decree 748/2019 for nedenstående (på knoldbasis):

- Sølvscurv (*Helminthosporium solani*): Max. 20% af knoldene, og max. 20% af overfladen
- Pulverscurv (*Spongospora subterranea*): Max. angrebsprocent 1%, max 10% af knoldoverfladen dækket
- Rodfiltsvamp (*Rhizoctonia solani*): Max. 5% med max. 10% af overfladen dækket
- Skurv (*Streptomyces scabies*): Max. 1.5 (Reference skurvskala 1.5 jf. NL-skala)
- Spirer: Max. 2 mm

Derudover må andre sygdomme/skader samlet set ikke overstige 5%:

- Fusarium: Max. 1%
- Vandsure (*Phytium* spp.): Max. 0,5%
- *Phoma* spp.: Max 0,5%
- Indre fejl/rust: Max. 0,5% (200 knolde gennemskæres)
- Mekaniske skader: Max. 3 %

Laboratorieundersøgelser

Sorten skal være testet hos pågældende avler og fundet fri for ring- og brunbakteriose.

Produktionsareal må ikke have været klassificeret som en nematodlokalitet inden for de seneste tre år.

Tillægserklæring på plantesundhedscertifikat

Import permit no. XXXXXXXXXX

The seed potatoes in this shipment are accepted as seed potatoes and the consignment is intended for cultivation only. The seed potatoes as well as their place of cultivation have been found free from:

- *Synchytrium endobioticum*
- *Leptinotarsa decemlineata*
- *Clavibacter michiganensis* subsp. *sepedonicus*
- *Ralstonia solanacearum*
- *Globodera rostochiensis*
- *Dickeya* spp.



The consignment was inspected and complies with Union Grade E certification scheme for virus tolerances of Union Grade E or Union Grade SE certification scheme for virus tolerances for seed potatoes of Union Grade SE. The general phytopathological conditions of seed potatoes according to the rules and regulations of the ARE (Decree No. 748 of 2019 of the Egyptian Ministry of Agriculture and annex no. 14 of the ministerial Decree no. 562/2019) are met.

Øvrigt



Forenede Arabiske Emirater

I henhold til: Conditions and Specifications of Imported Potato Tubers (mail fra 2014)

Marksyn

Partikontrol

Rhizoctonia solani: 10 % light Dutch Rhizoctonia scale 1

Dry-rot (incl. *Phoma* spp.): 0,50%

Wet-rot: 0%

External defects: 1%

Heavy internal defects: 1%

Sprouts caused by external pressure:

- < 2cm Ø and > 3mm deep max 6%
- 2cm Ø and > 3mm deep max 1%

Severe green color: 3%

Plant parasitic nematodes: 0%

Streptomyces scabies and *Spongospora subterranean*: did not exceed Dutchscab scale 1

Soil: max 05 %

Leptinotarsa decemlineata (Colorado beetle), *Globodera rostochiensis* (Potato cyst nematode), *Meloidogyne* spp. (Root knot nematodes), *Synchytrium endobioticum* (Potato wart disease), *Ralstonia solanacearum* (Brown rot), *Clavibacter michiganensis* subsp. *sepedonicus* (Ring rot)

Laboratorieundersøgelser

Partiet skal være testet for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

The tubers are free from *Leptinotarsa decemlineata* (Colorado beetle), *Globodera rostochiensis* (Potato cyst nematode), *Meloidogyne* spp. (Root knot nematodes), *Synchytrium endobioticum* (Potato wart disease), *Ralstonia solanacearum* (Brown rot), *Clavibacter michiganensis* subsp. *sepedonicus* (Ring rot) wet rot, plant parasitic nematodes, Broomrape (*Orobanche*) and Dodder (*Cuscuta*). *Rhizoctonia solani* did not exceed Dutch Rhizoctonia scale 10 % light.

Streptomyces scabies and *Spongospora subterranea* did not exceed Dutchscab scale 1. Dry rot did not exceed 0.5%. External defects and heavy internal defects did not exceed 1%. Severe green color did not exceed 3%. Medium internal defects did not exceed 6%. Spots < 2cm Ø and > 3mm deep did not exceed 6% and those > 2cm Ø and < 3mm deep caused by external pressure did not exceed 1%. Soil attached on tubers did not exceed 0.5% by weight.

Øvrigt



Irak – Model A

Model A: Uden Virusundersøgelse

Oplysninger fra: Technical regulations dealing with imported potato (modtaget af Solanum Nord 2002)

Marksyn

0% virus

Frihed for *Leptinotarsa decemlineata*

Partikontrol

Vådråd + tørråd: 1% (af knolde)

Pulverskurv: 0%

Dyb skurv: 0%

Overfladisk skurv >1/3 af overfladen: 5%

Pulverskurv: 0%

Rodfiltsvamp (sclerotier) > 1/20 af overfl: 0%

Rodfiltsvamp (sclerotier) < 1/20 af overfl: 10%

Jord (maks af vægt): 1%

Mekaniske skader og misdannelser: 1%

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være testet for ringbakteriose.

Tillægserklæring på plantesundhedscertifikat

The crops from which the seed potatoes originate were officially inspected whilst growing by inspectors of the Ministry, and were certified to conform to the standards of tolerance to virus diseases according to the Ministry's certification scheme. Potato yellowing vein 'virus', Potato yellow dwarf virus (PYDV), Potato Spindle Tuber Viroïd (PSTVd) has never been found in potatoes in Denmark. Tomato spotted wilt virus (TSWV) is known not to occur in this consignment. Potato tuber moth (*Phthorimaea operculella*), Japanese beetle (*Popillia japonica*) has never been found in Denmark. Colorado beetles (*Leptinotarsa decemlineata*) are known not to occur in the consignment. Sclerotium disease (*Sclerotium (Corticium) rolfii*), Dry rot (*Fusarium* spp.), Early blight (*Alternaria solani*) and Late blight (*Phytophthora infestans*) do not altogether exceed 1%.

The percentage of Common scab (*Streptomyces scabies*) is max 5 % of tubers showing lesions covering more than 1/3 of total tuber surface. The percentage of Powdery scab (*Spongospora subterranea*) is 0 %.

The percentage of Black scurf (*Rhizoctonia solani*) class E and A does not exceed 10% light. The seed potatoes originate from land which was officially sampled in [redacted] and examined for the presence of commercially harmful nematodes which would be subject to quarantine regulations. Soil samples were found to be free of the potato cyst nematodes *Globodera*



pallida and *Globodera rostochiensis*. The official inspection of the crops during the period of growth did not reveal any evidence of the presence of potato cyst nematodes.

The percentage of injured and irregular seed is less than 1%. Weight of soil does not exceed 1% of weight.

The consignment has been tested for and found free from Bacterial ring rot *Clavibacter michiganensis* subsp. *sepedonicus*.

Øvrigt



Irak – Model B

Model B: Med Virusundersøgelse

Oplysninger fra: Technical regulations dealing with imported potato (modtaget af Solanum Nord 2002)

Marksyn

Partikontrol

Vådråd + tørråd: 1% (af knolde)

Pulverskurv: 0%

Dyb skurv: 0%

Overfladisk skurv >1/3 af overfladen: 5%

Pulverskurv: 0%

Rodfiltsvamp (sclerotier) > 1/20 af overfl: 0%

Rodfiltsvamp (sclerotier) < 1/20 af overfl: 10%

Jord (maks af vægt): 1%

Mekaniske skader og misdannelser: 1%

Frihed for *Leptinotarsa decemlineata*

Laboratorieundersøgelser

Virusundersøgelse med følgende tolerancer:

- Præbasis: 1%
- Basis: 2%
- Certificeret Klasse A: 5%

Sorten fra den pågældende avler skal være testet for ringbakteriose.

Tillægserklæring på plantesundhedscertifikat

The crops from which the seed potatoes originate were officially inspected whilst growing, by Inspectors of the Ministry and were certified to conform to the standards of tolerance to virus diseases according to the Ministry's certification scheme. Potato yellowing vein 'virus', Potato yellow dwarf virus (PYDV), Potato Spindle Tuber Viroïd (PSTVd) has never been found in potatoes in Denmark. Tomato spotted wilt virus (TSWV) is known not to occur in this consignment.

Potato tuber moth (*Phthorimaea operculella*), Japanese beetle (*Popillia japonica*) has never been found in Denmark. Colorado beetles (*Leptinotarsa decemlineata*) are known not to occur in the consignment. Sclerotium disease (*Sclerotium (Corticium) rolfii*), Dry rot (*Fusarium* spp.), Early blight (*Alternaria solani*) and Late blight (*Phytophthora infestans*) do not altogether exceed 1%.

The percentage of Common scab (*Streptomyces scabies*) is max 5 % of tubers showing lesions covering more than 1/3 of total tuber surface.



The percentage of Powdery scab (*Spongospora subterranea*) is 0 %. The percentage of Black scurf (*Rhizoctonia solani*) class E and A does not exceed 10% light.

The seed potatoes originate from land which was officially sampled in _____ and examined for the presence of commercially harmful nematodes which would be subject to quarantine regulations. Soil samples were found to be free of the potato cyst nematodes *Globodera pallida* and *Globodera rostochiensis*.

The official inspection of the crops during the period of growth did not reveal any evidence of the presence of potato cyst nematodes. The percentage of injured and irregular seed is less than 1%. Weight of soil does not exceed 1% of weight. The consignment has been tested for and found free from Bacterial ring rot (*Clavibacter michiganensis* subsp. *sepedonicus*).

The consignment has passed through laboratory test for virus and meet the permissible percentages:

- Super Elite: less than 1%,
- Elite: less than 2%
- Class A: less than 5%

Øvrigt



Island

Information fra: Forordning om import og eksport af planter og planteprodukter (189/1990)

Marksyn

Produktionsarealet skal være fundet fri for coloradobiller (*Leptinotarsa decemlineata*).
Phytophthora infestans må ikke være observeret på kartoffelplanterne i vækstsæsonen.

Partikontrol

Knoldene må ikke have symptomer på *Phytophthora infestans*. Knoldene skal findes fri for *Fusarium* spp. og *Phoma exigua* var. *foveata*.

Laboratorieundersøgelser

Jordprøve skal udtages og analyseres for kartoffelcystenematoder (*Globodera pallida* og *Globodera rostochiensis*).

Knoldprøver skal udtages og analyseres for ringbakteriose (*Clavibacter michiganensis*) og brunbakteriose (*Ralstonia solanacearum*).

Tillægserklæring på plantesundhedscertifikat

Ingen.

Øvrigt

-



Israel

I henhold til: Requirements for import of seed potatoes from Denmark, 956/2023

Marksyn

Sortben / blødråd (*Pectobacterium* spp.), max 0,5%;

Sortben / blødråd (*Dickeya* spp.), max 0,03 %;

For kontrol af *Dickeya* og *Pectobacterium* kræves to marksyn, udført i løbet af vækstsæsonen; angrebne planter og deres knolde fjernes før høst.

Marksyn for frihed for Coloradobiller (*Leptinotarsa decemlineata*), som heller ikke må forekomme på pakkeriet.

Partikontrol

Kartoffelskurv (*Streptomyces* spp.):

- Max 30% og med max. 4 små pletter.
- Af de 30%: max 3% med mere end 5 pletter
- Max 0,3% med dyb skurv (over 3 mm)

Pulverskurv (*Spongospora subterranea*): Op til 1/8 af overfladen max 1%

Kartoffelskimmel (*Phytophthora infestans*): 0,30%

Rhizoctonia solani:

- Max 10% med op til 1/8 af overfladen inficeret
- Max 1% med mere end 1/8 af overfladen inficeret

Colletotrichum coccodes (black dots): Max 30% med op til 1/3 af overfladen inficeret

Virus Y^{ntm}: 0% symptomer på angreb

Tobacco rattle virus: 0% (én prøve pr mark - ved gennemskæring af knolde)

Helminthosporium solani (silver scurf): Max. 30 %

Meloidogyne spp.

Tolerance-procent er på antal og ikke vægt.

Laboratorieundersøgelser

Marken skal være testet for *Globodera pallida* og *Globodera rostochiensis* 0%

Partiet skal testes fri for brunbakteriose (*Ralstonia solanacearum*) (200 knolde pr. 25 ton)
0%

Tillægserklæring på plantesundhedscertifikat

The fields and the surrounding area (as specified below) have never been found to be infected with:

- A) *Synchytrium endobioticum* (potato wart disease) (2 km).
- B) *Ralstonia solanacearum* (brown rot disease) (5 km).
- C) *Clavibacter michiganensis* subsp. *sepedonicus* (potato ring rot disease) (10 km).



The fields from which the crop derived and the packing houses are free from *Leptinotarsa decemlineata* (Colorado potato beetle)

The fields from which the crop was derived were found free from *Globodera rostochiensis* (golden nematode) and *Globodera pallida* (white potato cyst nematode) according to official laboratory examinations of soil samples taken up to 10 months before planting or during the current potato seed crop growth close to harvest or directly after harvest of the current potato seed crop.

The fields from which the crop derived were inspected at least twice, during active growth, and the incidence of *Pectobacterium* spp. (formerly: *Erwinia carotovora carotovora* and *Erwinia carotovora atroseptica*) did not exceed 0.5%. All infected plants and their tubers were removed prior to harvest.

The fields from which the crop derived were inspected at least twice, during active growth and found free from *Pectobacterium carotovora* subsp. *brasilense* and *Candidatus phytoplasma solani* ((Stolbur phytoplasma).

The fields from which the crop derived were inspected at least twice, during active growth, and the incidence of *Dickeya* spp. (formerly: *Erwinia chrysanthemi* pathovars attacking *Solanaceae*) did not exceed 0.03%. All infected plants and their tubers were removed prior to harvest.

The country of production is free from Potato spindle tuber viroid in potatoes

Tuber Inspection/Testing (percentage of the consignment by tubers).

Tubers should be free of the following pests and diseases or not exceed the limit stated below:

Synchytrium endobioticum (potato wart disease) 0%

Ralstonia solanacearum (brown rot disease) based upon lab. Test of 200 tubers/25 ton 0 %

Dickeya solani (formerly: *Erwinia chrysanthemi*) 0%

Pectobacterium carotovora subsp. *brasilense* 0%

Candidatus phytoplasma solani (Stolbur phytoplasma) 0%

Clavibacter michiganensis subsp. *sepedonicus* (potato ring rot disease) 0%

Leptinotarsa decemlineata (Colorado potato beetle) 0%

Globodera rostochiensis (golden nematode) 0%

Globodera pallida (white potato cyst nematode) 0%

Ditylenchus destructor (potato rot nematode) 0%

Meloidogyne chitwoodi (Columbia root-knot nematode) 0%

PVY ntn 0%

Tobacco rattle virus 0%

Potato spindle tuber viroid 0%

Streptomyces spp. (common scab)

No more than 4 small spots 30%

More than 5 spots of each affected tuber as part of the 30% - maximum 3%

Deep pitted scab 0.3%



Spongospora subterranea (powdery scab) 1%

Severe cankerous form of the disease 0%

Phytophthora infestans (late blight) 0.3%

Soft rots caused by various organisms 0.1%

Fusarium spp. and *Phoma* spp. (dry rots) 1%

(Maximum cumulative tolerance for late blight, soft rots and dry rots mentioned above is 1%)

Rhizoctonia solani (black scurf) 10%

Colletotrichum coccodes (black dots) 30%

Helminthosporium solani (silver scurf) affected tuber 30%

Øvrigt

Israel har krav til både dyrknings- og omkringliggende arealer som følge;

Der må aldrig have været fundet kartoffelbrøk (*Synchytrium endobioticum*) indenfor 2 km.



Jordan – Model A

Oplysninger fra: Nederlandske og Skotske myndigheder

Med virusundersøgelse

Marksyn

Partikontrol

Visuelt syn af partiet kræves med:

- Vådforrådnelse 0 %
- Tørforrådnelse 1 %
- Rodfiltsvamp max 2 % (kraftigt angreb, hvor > 5 % af overfladen er dækket)
- Rodfiltsvamp max 25 % (svagt angreb, hvor < 5 % af overfladen er dækket)
- Pulverskurv 0 % (max 2-4 pletter pr. knold)
- Sølvskurv (svampet) max 1 % (må højst dække 5 % af overfladen)
- Meloidogyne spp.

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være undersøgt for ring- og brunbakteriose.

Partiet skal være undersøgt for virus (bladrulle, Y, A, X og S) ved virusundersøgelse med tolerancer til godkendelse som basisavl.

I forbindelse med udstedelse af plantesundhedscertifikat skal resultatet af virusundersøgelsen være dokumenteret i form af et fremvist analysebevis.

Tillægserklæring på plantesundhedscertifikat

The seed potatoes in this consignment have been officially inspected, tested and found free from pests and comply with the tolerances, as defined by 102, 103, 104, 105, 106, 107, 108, 109 et 110 of the Instruction No. Z/13 of the year 2022 (Instructions of Registration Conditions and Procedures of Vegetable Seeds and their Origins (Imported and Locally Produced) including their Production and Trading Conditions issued under Clause 17.B and 18.b of Agriculture Law No. 13 of 2015 (as amended)).

The seed potatoes were inspected during the whole vegetation period by the Danish Agricultural Agency

The seed potatoes were inspected prior to shipment.

The seed potatoes in this shipment were produced in a production site where the soil was officially tested and found free of nematodes *Globodera rostochiensis* and *Globodera pallida*.

The seed potatoes in this shipment were produced at an officially inspected production site and found free of *Popillia japonica*, *Tecia solanivora*, *Epitrix cucumeris*, *Epitrix similaris*, *Epitrix subcrinita*, *Epitrix tuberis*, *Candidatus Liberibacter solanacearum* et/and Pepino mosaic virus (PepMV).

The seed potatoes in this shipment have been officially inspected and/or tested and found free of *Ralstonia solanacearum* et/and *Clavibacter sepedonicus* (Syn. *Clavibacter michiganensis* subsp. *sepedonicus*). The seed potatoes originate in an area free from



Ralstonia solanacearum et/and *Clavibacter sepedonicus* (Syn. *Clavibacter michiganensis* subsp. *sepedonicus*).

The seed potatoes in this shipment have been officially inspected and/or tested and found free of *Meloidogyne chitwoodi*, *Meloidogyne fallax*, *Leptinotarsa decemlineata*, *Phthorimaea operculella*, *Ditylenchus destructor*, Potato spindle tuber viroid (PSTVd), Tobacco rattle virus (TRV), Potato mop-top virus (PMTV), Potato Virus S (PVS), *Fusarium oxysporum* f.sp. *albedinis*, *Fusarium foetens*, *Synchytrium endobioticum*, Candidatus *Phytoplasma asteris* (Syn. Aster yellows phytoplasma) et/and *Dickeya* spp.

The seed potatoes in this shipment have been officially inspected and/or tested and meet the allowable fungal disease infection levels defined by class for: *Phytophthora* spp (including *Phytophthora infestans*), *Alternaria* spp (including *Alternaria solani*), *Boeremia exigua* var. *exigua* (Syn. *Phoma exigua*), *Fusarium* spp, *Verticillium albo-atrum*, *Verticillium dahliae*, *Sclerotinia sclerotiorum* et/and *Colletotrichum caccodes* (anthracnose of potato).

The seed potatoes in this shipment have been officially inspected and/or tested and comply with the allowable bacterial disease infection rates defined by class for: *Pectobacterium atrosepticum* et/and *Pectobacterium carotovorum* (Syn *Pectobacterium carotovorum* subsp. *carotovorum*).

The seed potatoes in this shipment have been officially inspected and/or tested and comply with the allowable and class-defined virus infection levels for Potato leafroll virus (PLRV), Potato virus Y (PVY), Potato virus A (PVA), Potato virus X (PVX).

The seed potatoes in this consignment have been officially inspected and comply with the percentage of infection allowed and defined by class for: *Rhizoctonia solani*, *Streptomyces scabiei* (Syn. *Streptomyces scabies*), *Spongospora subterranea* et/and *Helminthosporium solani*.

Øvrigt



Jordan – Model B

Oplysninger fra: Nederlandske og Skotske myndigheder

Uden virusundersøgelse

Marksyn

0 % virus ved marksyn

Partikontrol

Visuelt syn af partiet kræves med:

- Vådforrådnelse 0 %
- Tørforrådnelse 1 %
- Rodfiltsvamp max 2 % (kraftigt angreb, hvor > 5 % af overfladen er dækket)
- Rodfiltsvamp max 25 % (svagt angreb, hvor < 5 % af overfladen er dækket)
- Pulverskurv 0 % (max 2-4 pletter pr. knold)
- Sølvscurv (svampet) max 1 % (må højst dække 5 % af overfladen)
- Meloidogyne spp.

Laboratorieundersøgelser

Partiet skal være undersøgt for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

The seed potatoes in this consignment have been officially inspected, tested and found free from pests and comply with the tolerances, as defined by 102, 103, 104, 105, 106, 107, 108, 109 et 110 of the Instruction No. Z/13 of the year 2022 (Instructions of Registration Conditions and Procedures of Vegetable Seeds and their Origins (Imported and Locally Produced) including their Production and Trading Conditions issued under Clause 17.B and 18.b of Agriculture Law No. 13 of 2015 (as amended)).

The seed potatoes were inspected during the whole vegetation period by the Danish Agricultural Agency

The seed potatoes were inspected prior to shipment.

The seed potatoes in this shipment were produced in a production site where the soil was officially tested and found free of nematodes *Globodera rostochiensis* and *Globodera pallida*.

The seed potatoes in this shipment were produced at an officially inspected production site and found free of *Popillia japonica*, *Tecia solanivora*, *Epitrix cucumeris*, *Epitrix similaris*, *Epitrix subcrinita*, *Epitrix tuberis*, *Candidatus Liberibacter solanacearum* et/and Pepino mosaic virus (PepMV).

The seed potatoes in this shipment have been officially inspected and/or tested and found free of *Ralstonia solanacearum* et/and *Clavibacter sepedonicus* (Syn. *Clavibacter michiganensis* subsp. *sepedonicus*). The seed potatoes originate in an area free from *Ralstonia solanacearum* et/and *Clavibacter sepedonicus* (Syn. *Clavibacter michiganensis* subsp. *sepedonicus*).

The seed potatoes in this shipment have been officially inspected and/or tested and found free of *Meloidogyne chitwoodi*, *Meloidogyne fallax*, *Leptinotarsa decemlineata*, *Phthorimaea operculella*, *Ditylenchus destructor*, Potato spindle tuber viroid (PSTVd),



Tobacco rattle virus (TRV), Potato mop-top virus (PMTV), Potato Virus S (PVS), *Fusarium oxysporum* f.sp. *albedinis*, *Fusarium foetens*, *Synchytrium endobioticum*, *Candidatus Phytoplasma asteris* (Syn. Aster yellows phytoplasma) et/and *Dickeya* spp.

The seed potatoes in this shipment have been officially inspected and/or tested and meet the allowable fungal disease infection levels defined by class for: *Phytophthora* spp (including *Phytophthora infestans*), *Alternaria* spp (including *Alternaria solani*), *Boeremia exigua* var. *exigua* (Syn. *Phoma exigua*), *Fusarium* spp, *Verticillium albo-atrum*, *Verticillium dahliae*, *Sclerotinia sclerotiorum* et/and *Colletotrichum caccodes* (anthracnose of potato).

The seed potatoes in this shipment have been officially inspected and/or tested and comply with the allowable bacterial disease infection rates defined by class for: *Pectobacterium atrosepticum* et/and *Pectobacterium carotovorum* (Syn *Pectobacterium carotovorum* subsp. *carotovorum*).

The seed potatoes in this shipment have been officially inspected and/or tested and comply with the allowable and class-defined virus infection levels for Potato leafroll virus (PLRV), Potato virus Y (PVY), Potato virus A (PVA), Potato virus X (PVX).

The seed potatoes in this consignment have been officially inspected and comply with the percentage of infection allowed and defined by class for: *Rhizoctonia solani*, *Streptomyces scabiei* (Syn. *Streptomyces scabies*), *Spongospora subterranea* et/and *Helminthosporium solani*.

Øvrigt



Kanariske Øer (Spanien)

Kontakt Landbrugs- og Fiskeristyrelsen for eksportbetingelser for kartofler til De Kanariske Øer: Sundhedfroel@lfst.dk



Kosovo

Oplysninger fra NL hjemmeside

Marksyn

Partikontrol

Laboratorieundersøgelser

Frihed for *Globoedra pallida* og *Globodera rostochiensis*.

Frihed for *Clavibacter sepedonicus* og *Ralstonia solanacearum* testet på sorten hos avleren.

Tillægserklæring på plantesundhedscertifikat

The place of production has been found free from A1 and A2 quarantine pests. *Synchytrium endobioticum* does not occur within a radius of 10 km. from the place of production. The place of production from which the seed potatoes originate is free from *Clavibacter michiganensis* subsp. *sepedonicus*.

Prior to planting the soil of the fields of production has been tested and found free from *Globodera pallida* and *Globodera rostochiensis*. The seed potatoes were inspected during the vegetation period and were found free from virus to an extent ensuring certification as class SE, E and/or A.

The consignment was inspected and found free from *Clavibacter michiganensis* subsp. *sepedonicus*, *Pseudomonas (Ralstonia) solanacearum*, potato spindle tuber pospiviroid and other non-European viruses and virus like organisms.

Soil has been inspected and found free from presence of quarantine species of nematodes.

The consignment was inspected by the competent Plant Protection Organization of Denmark.

Øvrigt



La Réunion (Frankrig)

Oplysninger hentet fra: "ARRÊTE PRÉFECTORAL N° 2011 – 1479 du 30 septembre 2011 modifié, conditions requises pour l'introduction de végétaux et produits végétaux à la Réunion." and "ANNEXE IV exigences particulières requises pour l'introduction et la circulation des végétaux et produits végétaux et autres objets sur le territoire de l'île de la Réunion"

Marksyn

Partikontrol

Partiet skal synes for *Epitrix* sp., *Meloidogyne chitwoodi* og *Meloidogyne fallax*, *Tecia solanivora*, *Synchytrium endobioticum* og Potato spindle tuber viroid.

Laboratorieundersøgelser

Analysebeviser for *Clavibacter sepedonicus* og *Ralstonia solanacearum*.

Tillægserklæring på plantesundhedscertifikat

According to "ARRÊTE PRÉFECTORAL N° 2011 – 1479 du 30 septembre 2011 modifié, conditions requises pour l'introduction de végétaux et produits végétaux à la Réunion." and "ANNEXE IV exigences particulières requises pour l'introduction et la circulation des végétaux et produits végétaux et autres objets sur le territoire de l'île de la Réunion"

9 a) and b) the consignment has been inspected and/or tested free from harmful organisms before shipment.

57 a) the potato tubers come from an area known to be free from *Synchytrium endobioticum*.

57.2 a) the potato tubers come from an area known to be free from *Epitrix* sp. (Angiv region)

58 a) the potato tubers come from an area known to be free from *Clavibacter michiganensis* spp.

59 a) the potato tubers come from an area known to be free from Potato spindle tuber viroid.

60 the potato tubers come from a field tested free from *Globodera rostochiensis* and *Globodera pallida*

60 aa) the potato tubers come from an area known to be free from *Ralstonia solanacearum*.

60 cc) the potato tubers come from an area where *Meloidogyne chitwoodi* and *Meloidogyne fallax* is not known to occur

62 a) the potato tubers come from an area where *Tecia solanivora* is not known to occur

63 no symptoms of stolbur phytoplasma have been observed on the place of production since the last complete vegetation cycle.

Øvrigt



Libanon

Decree No. 693/1 conditions to import potato seeds for the 2016/17 season.

NB Krav om læggekartofler af klasse E eller højere (SE). Virustolerancen kan enten opfyldes ved laboratorietest eller marksyn.

Fysisk kontrol af partiet: (NB: tolerance-procent er på antal og ikke vægt)

Marksyn

(*Pectobacterium*, *Dickeya* spp.) 0% sortben

Klasse E max. 1,0% virus.

Klasse SE: max 0,75% virus.

Partikontrol

- Blødråd: 0%
- Tørråd*: 0,5%
- *Phoma*: 0%
- Sølvscurv: ingen rynkede knolde
- Spirer: max. 15 mm på 20% af knoldene
- *Ditylenchus dipsaci* – tørråd: 0%
- *Rhizoctonia* og *Polyscytalum*: max. 5% med infektion (antal), og inficeret område må ikke overstige 10% af overfladen
- Kartoffel- og pulverscurv: 33% af knoldene (antal) skal være fri for infektion af alm. scurv og pulverscurv, og i de resterende 66% af knoldene må der af de inficerede ikke være mere end 2-4 svage symptomer som ikke må dække mere end 1/16 del af overfladen
- Jord: Max. 1%
- Overfladeskader (mekaniske og insekt): Max. 2%
- *Meloidogyne* spp.

*) "Tørråd" omfatter *Phytophthora infestans*, *Alternaria solani*, *Fusarium solani*, *Verticillium* spp.

Behandling: Fungicidbehandling kræves.

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være testet for ring- og brunbakteriose

Tillægserklæring på plantesundhedscertifikat

The seed potatoes originate from an area where *Synchytrium endobioticum* (Schilb.) Perc. and *Ralstonia (Pseudomonas) solanacearum* is not known to occur for at least five years and from *Clavibacter michiganensis* ssp. *sepedonicum* for at least three years.

The potatoes are absolutely free from Colorado potato beetle (*Leptinotarsa decemlineata*), Potato tuber moth (*Phthorimaea (Gnorimoschima) operculella*), Japanese beetle (*Popillia*



japonica), Sugar beet weevil (*Asproparthenis punctiventris*), Potato Rot Nematode (*Ditylenchus destructor*), Stem and bulb nematode (*Ditylenchus dipsaci*), White potato cyst nematode (*Globodera pallida*), Golden nematode (*Globodera rostochiensis*), Potato gangrene (*Phoma foveata*), Tomato foot rot (*Phytophthora cryptogea*), Pink tuber rot (*Phytophthora erythroseptica* var. *erythroseptica*), Wart disease of potato (*Synchytrium endobioticum*), Corm scab (*Burkholderia gladioli* pv. *gladioli*), potato ring rot (*Clavibacter michiganensis* ssp. *sepedonicus*), Bacterial wilt of Dahlia (*Erwinia chrysanthemi* pv. *chrysanthemi*), Black leg (*Pectobacterium* (*Erwinia*) *carotovora* ssp. *atroseptica*), Soft rot (*Pectobacterium* (*Erwinia*) *carotovora* ssp. *carotovora*), Lettuce marginal leaf blight (*Pseudomonas marginalis* pv. *marginalis*), Brown rot (*Pseudomonas solanacearum*), and Potato spindle tuber viroid (PSTVd).

The seed potatoes are substantially free from Late blight (*Phytophthora infestans*), Early blight (*Alternaria solani*), *Fusarium* dry rot (*Fusarium solani*) and *Verticillium* (*Verticillium* sp.), less than 0,5 %. The consignment is free from wrinkled tubers due to silver scurf (*Helminthosporium solani*).

Not more than 5% of the tubers (by number) are infected with Skin spot (*Oospora pustulans*) or Black scurf (*Rhizoctonia solani*), and the infected area of the tuber does not exceed 10 %.

33 % of the tubers (by number) are not infected with Common scab (*Streptomyces scabies*) and Powdery scab (*Spongospora subterranea*), and the tubers infected in the 66% do not exceed 4-2 slight wounds covering an area less than 1/16 of the tuber area.

Not more than 1 % of potato seeds Class E or 0,75 % of potato seeds Class SE are attacked by viral diseases.

The consignment has been inspected and found substantially free from *Meloidogyne chitwoodi* and *Meloidogyne fallax*.

The Danish Agricultural Agency hereby confirm that the goods were inspected by the Danish Agricultural Agency, and that the norms and standards of this consignment comply with the clauses mentioned in the Lebanese Ministerial Decree No. 693/1 concerning the conditions for potato seeds which can be exported to Lebanon as per minagri [redacted] tender book.

Øvrigt



Libyen

I henhold til: The State of Libya The National Concord Government Ministry of Agriculture,
No. 17/2019 (23/01/2019)

Marksyn

Potato leaf roll virus: 0,1%

Potato mosaic virus: 0,1%

Leptinotarsa decemlineata (Coloradobille) og *Popillia japonica* (Japanbille): 0%

Pectobacterium carotovorum subsp. *Carotovorum*: 0%

Partikontrol

- Almindelig skurv max 25% med to ikke sammenhængende pletter eller små pletter på max. 1/32 del af knolde
- *Clavibacter michiganensis* subsp. *Sepedonicus*: 0%
- *Fusarium* (dry rot): Max. 0,5%
- *Helminthosporium solani* (sølvskurv): max. 10% af knoldene og max. 10% af overfladen
- *Phytophthora infestans*: max. 0,5%
- *Ralstonia solanacearum*: 0%
- *Rhizopus oryzae* (soft rot disease): 0%
- *Sclerotinia sclerotiorum*: Max. 0,5%
- *Spongospora subterranea*: (pulverskurv): 0%
- *Thanatephorus cucumeris*: Max. 5% af knoldene og max. 10% af overfladen
- Spirer: praktisk taget fri
- Jord: praktisk taget fri
- Mekaniske- og fysiologiske skader: Max. 0,5% af vægten
- *Synchytrium endobioticum*: 0%
- *Meloidogyne* spp.

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være testet for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

The seed potatoes were grown in land in which Potato Wart Disease (*Synchytrium endobioticum*) has been known not to occur at any time nor within 10 kilometres thereof.

The seed potatoes originate from land which was officially sampled in [redacted] and examined for the presence of commercially harmful nematodes which would be subject to quarantine regulations. Soil samples were found to be free of the potato cyst nematodes *Globodera pallida* and *Globodera rostochiensis*. The official inspection of the crops during the period of growth did not reveal any evidence of the presence of potato cyst nematodes.

Potato tuber moth (*Phthorimaea operculella*), Japanese beetle (*Popillia japonica*), Potato brown rot (*Ralstonia solanacearum*), Potato Stolbur phytoplasma, Potato yellowing vein 'virus', Potato yellow dwarf virus (PYDV) and Potato Spindle Tuber Viroïd (PSTVd) has never been found in potato in Denmark.



Colorado beetles (*Leptinotarsa decemlineata*), Potato rot nematode (*Ditylenchus destructor*), Potato ring rot (*Clavibacter michiganensis* subsp. *sepedonicus*) and Tomato spotted wilt virus (TSWV) are not known to occur in the consignment.

The consignment has been inspected and found substantially free from *Meloidogyne chitwoodi* and *Meloidogyne fallax*.

The crops from which the seed potatoes originate were officially inspected whilst growing, by Inspectors of the Ministry and were certified to conform to the standards of freedom of virus diseases according to the Ministry's certification scheme.

Øvrigt



Marokko

Information fra: no. 593-17 of 15 kaada 1438 (8. august 2017)

Marksyn

Partikontrol

- *Phytophthora infestans* 2 %
- Blødråd 2 %
- Sølvscurv (*Helminthosporium solani*) og alm. skurv: max. 5 % af knoldene med symptomer på mere end 1/3 af overfladen
- Rodfiltsvamp: max. 10 % af knoldene (antal)
- Frihed for *Leptinotarsa decemlineata*
- Praktisk taget fri for jord og halm (Frihed for jord og halm jf. Marokkos lovgivning)
- *Meloidogyne* sp.

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være testet for ringbakteriose.

Tillægserklæring på plantesundhedscertifikat

The shipment is tested and/or inspected and found free of:

Thecaphora solani, *Ralstonia solanacearum*, *Clavibacter michiganensis* subsp. *sepedonicus*, *Clavibacter michiganensis* subsp. *michiganense*, *Leptinotarsa decemlineata*, *Synchytrium endobioticum*, *Spongospora subterranea*, *Angiosorus solani*, Potato spindle tuber viroid, Potato mop-top virus, *Liberibacter solanacearum*, *Dickeya dianthicola*, *Dickeya chrysanthemi*, *Globodera rostochiensis*, *Globodera pallida*, *Xanthomonas campestris* pv. *versicatoria*, *Meloidogyne* spp. et *Ditylenchus destructor*.

The shipment is free of:

Cirsium arvense, *Cuscuta* spp. et *Orobanche* spp.

Øvrigt



Montenegro

Information fra: Rulebook of Montenegro List IVA part I from 14 December 2023

Marksyn

Partikontrol

Laboratorieundersøgelser

Partiet skal være testet for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

The tubers in this consignment comply with Law on Plant health Protection in Montenegro List IVA part I from 14 December 2023 (p.1739) harmonized with (EU) 2019/2072. Tubers comply with (14) that the consignment or lot does not contain more than 1% by net weight soil. (15.a) *Tecia solanivora* does not occur in Denmark. (16.a) Denmark is known to be free from *Clavibacter sepedonicus*. (17.a) The tubers originate in areas known to be free from *Synchytrium endobioticum*, and no symptoms have been observed either at the place of production or it's immediate vicinity. (18) The tubers originate from a field known to be free from *Globodera rostochiensis* and *Globodera pallida* with appropriate methods. (19.b) The tubers originate from a place of production found free from *Ralstonia solanacearum*, *Ralstonia pseudosolanacearum*, *Ralstonia syzygii* subsp. *celebensis* and *Ralstonia syzygii* subsp. *indonesiensis* and are subjected to official testing after harvest. (20.a or 20.d) Depending on the area of production tubers originate from areas where *Meloidogyne chitwoodi* and *Meloidogyne fallax* are known not to occur or the tubers have been checked for the presence of symptoms after an appropriate method to induce symptoms, or laboratory tested, as well as inspected visually both externally and by cutting the tubers.

Øvrigt



Nordmakedonien

Oplysninger fra: Book of rules on lists of harmful organisms, plants, plant products and other objects and items

Marksyn

Partikontrol

Laboratorieundersøgelser

Frihed for *Globodera pallida*, *Globodera rostochiensis*

Tillægserklæring på plantesundhedscertifikat

The consignment complies with Rule 91/21 Annex 4 - List IV Part A – section 1 item 25.2 (a) the tubers originate in countries recognized as free from *Clavibacter michiganensis* subsp. *sepedonicus* (Spieckermann and Kotthoff) Davis et al., 25.4 the tubers originate in an area known to be free of *Globodera rostochiensis* (Wollenweber) Behrens and *Globodera pallida* (Stone) Behrens., 25.4 (aa) the tubers originate in areas where *Ralstonia solanacearum* (Smith) Yabuuchi et al is known not to occur, 25.4 (cc) the tubers originate in areas where *Meloidogyne chitwoodi* Golden et al. (all populations) and *Meloidogyne fallax* Karssen are known not to occur and 25.4.2 (a) the tubers originate in a country where *Scrobipalopsis solanivora* Povolny is known not to occur

Øvrigt

Quality certificate skal vedlægges plantesundhedscertifikatet



Saudi Arabien – Model A

(Krav fra: Ministerial Decision No. 246999 dated 20/08/1431 H og 4/3/49943, dated 9/7/1415 H)

Model A – UDEN krav om lab.-analyse for virus Y, A, X, S, bladrullevirus

Krav til import af læggekartofler:

Import er kun tilladt for klasse E eller højere klasser. Størrelse: 28-55 mm

Marksyn

- Sortben (*Pectobacterium* spp.) 0 %
- Virus 0 %
- *Verticillium dahliae*, *Verticillium albo-atrum* 0,1 %

Partikontrol

Tolerancen angivet i pct. knolde

- Blødråd, sortben (arter af *Dickeya*, *Pectobacterium*): 0 %
- Ringrust; TRV og moptopvirus: 0 %
- Tørråd (*Phoma*, kartoffelskimmel, *Alternaria*): 0,5 %
- Fusarium: 0 %
- *Rhizoctonia*: 0 % svær infektion (> 1/20 af overfladen)
- *Rhizoctonia*: max 10 % let infektion (<1/20 af overfladen)
- Alm. skurv (*Streptomyces*): 5 %, max. 1/3 af overfladen
- Pulverskuv (*Spongospora*): 3 %, max. 1/20 af overfladen
- Sølvskurv og indskrumpne knolde: 1 %; skurv max. 1/20 af overfladen
- Spirer: 1 % må have spirer længere end 10 mm,
- Grønne knolde, beskadigelser, hollow heart, brown/black heart, i alt: max 3 %
- Jord og sand i hver sæk: 0,5 % (men knolde helt frie for jord)
- Fungicidbehandling (Imazalil) af hele partiet
- Mærkningskrav (ud over alm plantepas): Importørens navn og adresse

Laboratorieundersøgelser

Sorten fra den pågældende avler skal testes fri for ring- og brunbakteriose (CMS og RS)

Tillægserklæring på plantesundhedscertifikat

The seed potatoes are free from bacterial, and fungal diseases and free from insect pests, nematodes and soil according to Ministerial Decision No. 246999 dated 20/08/1431 H and Decision no. 4/3/49943, dated 9/7/1415 H and totally free 100% from brown rot disease caused by *Pseudomonas (Ralstonia) solanacearum*.

Øvrigt

Risø-erklæring, samt Non-GMO-erklæringen

Både Risøerklæringen og frihedserklæringen for GMO findes på lfst.dk



Saudi Arabien – Model B

(Krav fra Ministerial Decision No. 246999 dated 20/08/1431 H og 4/3/49943, dated 9/7/1413 H)

Model B med krav om laboratorieanalyse for virus

Krav til import af læggekartofler:

Import er kun tilladt for klasse E eller højere klasser. Størrelse: 28-55 mm

Marksyn

- Sortben (*Pectobacterium* spp.): 0 %
- *Verticillium dahliae*, *Verticillium albo-atrum*: 0,1 %

Partikontrol

Tolerancen angivet i pct. knolde

- Blødråd, sortben (arter af *Dickeya*, *Pectobacterium*): 0 %
- Ringrust; TRV og moptopvirus: 0 %
- Tørråd (*Phoma*, kartoffelskimmel, *Alternaria*): 0,5 %
- *Fusarium*: 0 %
- *Rhizoctonia*: 0 % svær infektion (> 1/20 af overfladen)
- *Rhizoctonia*: max 10 % let infektion (<1/20 af overfladen)
- Alm. skurv (*Streptomyces*): 5 %, max. 1/3 af overfladen
- Pulverskuv (*Spongospora*): 3 %, max. 1/20 af overfladen
- Sølvskurv og indskrumpne knolde: 1 % skurv max 1/20 af overfladen
- Spirer: 1 % (må have spirer længere end 10 mm)
- Grønne knolde, beskadigelser, hollow heart, brown/black heart, i alt: max 3 %
- Jord og sand i hver sæk: 0,5 % (men knolde helt frie for jord)
- Fungicidbehandling (Imazalil) af hele partiet
- Mærkningskrav (ud over alm plantepas): Importørens navn og adresse

Laboratorieundersøgelser

- Ring- og brunbakteriose CMS og RS (sorten fra den pågældende avler)
- Virus: Bladrulle, Virus Y, Virus A: 2 %
- Virus: Virus X, Virus S: 4 %

Tillægserklæring på plantesundhedscertifikat

The seed potatoes are free from bacterial, and fungal diseases and free from insect pests, nematodes and soil according to Ministerial Decision No. 246999 dated 20/08/1431 H and Decision no. 4/3/49943, dated 9/7/1415 H and totally free 100% from brown rot disease caused by *Pseudomonas (Ralstonia) solanacearum*. We hereby state that the ratios of virus diseases in the seed potatoes are according to Saudi Arabian Ministry of Agriculture & Water Ministerial Decision No 246999, dated 20/8/1431H.

Øvrigt

Risø-erklæring, samt Non-GMO-erklæring

[Både Risøerklæringen og frihedserklæring for GMO findes på lbst.dk](http://lbst.dk)



Serbien

Oplysninger fra: Rulebook on lists of harmful organisms and lists of plants, plant products and regulated objects (7/2021, 22/2012, 57/2015).

Marksyn

Partikontrol

Laboratorieundersøgelser

Partiet skal være testet fri for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

Uden partikontrol:

The consignment complies with Law on Plant Health no. 7/2010, 22/2012 and 57/2015 List IV A part I 25.1 (a), 25.2 (a), 25.4 (a), (b), (c) and (d) and 25.4.2 (a).

Soil has been tested and found free from *Globodera rostochiensis* (Wollenweber) Behrens and *Globodera pallida* (Stone) Behrens.

Øvrigt



Syrien

I henhold til: Phytosanitary Requirements for Seeds potato Importation into Syria for the Season 2024/2025

The Syrian Arab Republic, Ministry of Agriculture and Agrarian Reform, Directorate of Plant Protection

Marksyn

Partikontrol

Kontrollen skal omfatte et minimum på 2 – 5% af sækkene

- Tørråd (skimmel, *Phoma* spp., *Phytophthora infestans*), deforme knolde og mekaniske skader: max. 1 % knolde
- Blødråd: max. 1 knold pr. 3 sække (150 kg)
- Sølvskurv (tør og rynket): max. 1 knold pr. sæk (50 kg)
- Sølvskurv (sølvgrå pletter): max. 40 % af overfladen
- Skurv (alle typer): max. 10% af knoldene: højst 25% af knoldoverfladen inficeret, resterende 90%: højst 5% af knoldoverfladen inficeret
- Pulverskurv, tenknold, alle insekter; 0 %
- Jord i forsendelsen: max. 100 gr. pr. 50 kg, knoldene fri for jord
- Rodfiltsvamp (*Rhizoctonia solani*): max. 5% af knoldene: højst 5% af knoldoverfladen inficeret, resterende 95%: højst 1% af knoldoverfladen inficeret
- Knolde med spirer > 2 mm, max. 2 % knolde
- Frostskeer: 0 %
- *Leptinotarsa decemlineata*; Produktionsarealet skal være frit

Laboratorieundersøgelser

Sorten hos den pågældende avler skal være testet for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

The seed potatoes were treated with suitable materials that protect seed potatoes and prevent fungal diseases and the seed potatoes are not genetically modified.

The seed potatoes were officially tested in Denmark for freedom of BROWN ROT and RING ROT. PSTVd is not known to exist on potatoes in Denmark. They were inspected in Denmark and found free from viral diseases as well as other diseases according to the specifications set out in article 3 of No94/T date 1/8/2024 "Phytosanitary Requirements for Seeds potato Importation into Syria for the season 2024-2025" issued by the Plant Protection Directorate, Ministry of Agriculture and Agrarian Reform, The Syrian Arab Republic. The wood packaging materials have been treated according to the ISPM No.15

Øvrigt

Risø-erklæring og frihedserklæring for GMO

[Både Risøerklæringen og frihedserklæring for GMO findes på lbst.dk](https://lbst.dk)

Træemballage skal være certificeret ISPM 15.

Knolde skal være behandlede med fungicid, og pakket i nye sække.



Tunesien

Oplysninger fra Cahier des charges, Importation de semences de pommes de terre – campagne 2017-2018

Marksyn

Partikontrol

(grænseværdier angivet i vægtprocent)

Alle skal være certificerede til mindst klasse A

1. Jord og andre fremmedlegemer: 0,5%
 2. Deforme eller beskadigede knolde: 1 %
 3. Våd råd (*Erwinia* spp., *Phytophthora erythroseptica*, *Pythium* sp) og skimmel (*Phytophthora infestans*): 0,1%
 4. Tørråd (*Fusarium* spp, *Verticillium* spp & *Phoma* spp. undtagen *Phoma andina* og *Phoma exigua* var. *foveata*): 0,2%
 5. Smælder, kartoffelmøl, orme og snegle: 3%
 6. Pkt. 1.-5. tilsammen max 4%
 7. Alm. Skurv: 10% angrebsgrad 3 (se ill.*)
 8. Sølvskurv: Med let angreb (3-4 pletter/knold): max 10%
 9. Rodfiltsvamp: max. 10% med angrebsgrad 3 og max 5 % med angrebsgrad 3 - 7 (se ill.*)
- Pkt. 6.-7. tilsammen max 10%

10. Spiring må ikke forekomme i mere end 10% af kartoflerne

* se 2-Attest Tunesien Skurv + Rodfiltsvamp skala.jpg

Laboratorieundersøgelser

Sorten hos avler skal være undersøgt for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

The consignment was officially inspected before shipment and found free from harmful pests according to Cahier des charges, Importation de semences de pommes de terre – campagne 2017-2018.

The seed potatoes originate in an area where *Clavibacter michiganensis* subsp. *sepedonicus*, *Ralstonia solanacearum*, *Synchytrium endobioticum*, *Globodera pallida*, *Globodera rostochiensis*, Stolbur phytoplasma and Potato spindle tuber viroid is not known to occur.

Øvrigt



Tyrkiet

Oplysninger fra: Annex I, II and IV til Regulation om Plant Quarantine

Marksyn

Partikontrol

Phoma: 0%

Pulverskurv: 0% knolde med åbne pustler

Meloidogyne spp: 0%

Ditylenchys dipsaci: 0%

Laboratorieundersøgelser

Sorten fra den pågældende avler skal være testet for ring- og brunbakteriose.

Tillægserklæring på plantesundhedscertifikat

The consignment is free from harmful organisms according to Regulation on Plant Quarantine Annex I and II and comply with requirements in Annex 4 point 32.1, 32.2 a), 32.3, 32.4 a), b), c), d) and e), 32.4.1 and 32.4.2 a) and 32.6.

Øvrigt

Seed certificate, Türkiye



UK – Meristem

Information hentet fra: Great Britain's Phytosanitary Conditions Regulation (2019/2072), Annex 7

Marksyn

Partikontrol

Ingen partikontrol, men skal anmeldes til fysisk eksportkontrol.

Laboratorieundersøgelser

Ingen

Tillægserklæring på plantesundhedscertifikat

This consignment complies with, item 9 a (ii) of Annex 7 to Great Britain's Phytosanitary Conditions Regulation (2019/2072) in that no symptoms of Tomato leaf curl New Delhi Virus have been observed on the plants during their complete cycle of vegetation, and item 28 b that no symptoms of *Candidatus Phytoplasma 'solani'* Quaglino et al. have been observed on the plants at the place of production since the beginning of the last complete cycle of vegetation.

Øvrigt



UK – Eksport til avler

Information hentet fra: The Plant Health (Phytosanitary Conditions) (Amendment) (EU Exit) Regulations 2020, Schedule 7

Marksyn

Partikontrol

Laboratorieanalyser

Tillægserklæring på plantesundhedscertifikat

The shipment complies with the phytosanitary requirements of item 19 (a) of Annex VII to Great Britain's Phytosanitary Conditions Regulation (2019/2072). The tubers originated in a country where *Tecia solanivora* is known not to occur

The shipment complies with the phytosanitary requirements of item 20 (a) i of Annex VII to Great Britain's Phytosanitary Conditions Regulation (2019/2072). The potato tubers have originated in an area known to be free from *Clavibacter sepedonicus* (Spieckermann & Kotthoff) Li et al.

The shipment complies with the phytosanitary requirements of item 20 (b) of Annex VII to Great Britain's Phytosanitary Conditions Regulation (2019/2072). The potato tubers have originated in a place of production free from *Synchytrium endobioticum* (Schilbersky Percival).

The shipment complies with the phytosanitary requirements of item 20 (c) i of Annex VII to Great Britain's Phytosanitary Conditions Regulation (2019/2072). The potato tubers have originated in an area known to be free from *Ralstonia solanacearum* (Smith) Yabuuchi et al. emend. Safni et al.

The shipment complies with the phytosanitary requirements of item 20 (d) i of Annex VII to Great Britain's Phytosanitary Conditions Regulation (2019/2072). The potato tubers come from an area where *Meloidogyne chitwoodi* Golden et al. is known not to occur.

The shipment complies with the phytosanitary requirements of point 20 (e) of Annex VII to Great Britain's Phytosanitary Conditions Regulation (2019/2072). The potato tubers come from a production site where the procedures for the control of *Globodera pallida* (Stone) Behrens and *Globodera rostochiensis* (Wollenweber) Behrens set out in EPPO PM 9/26 have been implemented.

The shipment complies with the phytosanitary requirements of point 28 (b) of Annex VII to Great Britain's Phytosanitary Conditions Regulation (2019/2072). No symptoms of *Candidatus Phytoplasma solani* Quaglino et al. have been observed on the plants at the place of production since the beginning of the last complete cycle of vegetation

Øvrigt



Ukraine

Krav hentet fra Nederlandene og Skotland

Marksyn

Partikontrol

Laboratorieundersøgelser

Globodera pallida, *Globodera rostochiensis*

Tillægserklæring på plantesundhedscertifikat

The seed potatoes were grown in land in which Potato Wart Disease (*Synchytrium endobioticum*) has not been known to occur at any time.

The seed potatoes originate from land which was officially sampled and examined for the presence of commercially harmful nematodes which would be subject to quarantine regulations. Soil samples were found to be free of the potato cyst nematodes *Globodera pallida* and *Globodera rostochiensis*. The official inspection of the crops during the period of growth did not reveal any evidence of the presence of potato cyst nematodes.

The official inspection of the crops during the period of growth did not reveal any evidence of the presence of Colorado beetles (*Leptinotarsa decemlineata*), Potato rot nematode (*Ditylenchus destructor*), Potato ring rot (*Clavibacter michiganensis* subsp. *sepedonicus*) and Tomato spotted wilt virus (TSWV).

Potato tuber moth (*Phthorimaea operculella*), Japanese beetle (*Popillia japonica*), Potato brown rot (*Ralstonia solanacearum*), Potato Stolbur phytoplasma, Potato yellowing vein 'virus', Potato yellow dwarf virus (PYDV) and Potato Spindle Tuber Viroïd (PSTVd) has never been found in Denmark on potatoes.

The tubers come from an area free from *Meloidogyne chitwoodi* and *Meloidogyne fallax*.

The crops from which the seed potatoes originate were officially inspected whilst growing, by Inspectors of the Ministry and are certified to conform to the standards of freedom of virus diseases according to the Ministry's certification scheme.

Øvrigt